

Canadian guidelines for body weight classification in adults: application in clinical practice to screen for overweight and obesity and to assess disease risk

Since 1985, the number of obese adults in Canada has doubled.¹ According to 2003 data from Statistics Canada (www.statcan.ca/english/freepub/82-221-XIE/00604/tables/html/1228_03.htm), 15% of adults are obese (body mass index [BMI] > 30 kg/m²) and 33% are overweight (BMI 25–29.9 kg/m²) and at increased risk of becoming obese. Given the alarming increase in the prevalence and potential health consequences of obesity, screening for overweight and obesity should be considered an integral part of a routine health assessment.

In 2003, Health Canada published the Canadian Guidelines for Body Weight Classification in Adults (available at www.hc-sc.gc.ca/hpfb-dgpsa/onpp-bppn/cg_bwc_introduction_e.html) to provide a framework for screening people for overweight and obesity. In this article we outline how this new weight classification can be applied to adult men and women (nonpregnant and nonlactating) between the ages of 18 and 65.

Why is screening important?

Obese people are at increased risk of diseases such as type 2 diabetes mellitus, hypertension, dyslipidemia, coronary artery disease, stroke, sleep apnea, osteoarthritis, gallstones, stress incontinence, depression and certain types of cancer.

Screening for overweight and obesity may change patient care in 3 ways. First, in adults who are found to be obese and also have obesity-related diseases, modest weight loss (5%–10% of total body weight) may improve control of these diseases. For example, weight loss in obese adults with type 2 diabetes and hypertension may improve glycemic and blood pressure

control and reduce drug therapy requirements. Second, in adults who are found to be obese but do not have obesity-related diseases, lifestyle interventions (e.g., starting an exercise program) may reduce the risk of these diseases. Third, in adults who are found to be overweight but are otherwise healthy, promoting healthy lifestyle practices may prevent them from becoming obese.

What are the methods for screening?

BMI and waist circumference are the 2 measures currently used to screen for overweight and obesity and to assess a person's risk of health problems (Table 1). The BMI is strongly correlated with reference measures of body fat (e.g., underwater weighing) and is an indicator of risk of weight-related diseases in men and women. The waist circumference measures abdominal (central) body fat and is strongly correlated with risk of health problems (type 2 diabetes, hypertension, dyslipidemia and the metabolic syndrome, which combines all 3 of these conditions).

What are some practical considerations?

Measuring the BMI and waist circumference: The BMI and waist circumference are simple, easy-to-use measures of adiposity. The correct way to measure the BMI is shown in Fig. 1, which also provides a BMI reference chart. The correct way to measure waist circumference is shown in Fig. 2.

Considering both measurements to assess disease risk: Although the BMI and waist circumference are related, the latter provides an in-

dependent estimate of health risk beyond that determined by the BMI. It is particularly useful to consider both measures when assessing a patient's disease risk (Fig. 3). For example, in a person with a normal BMI, an increased waist circumference may indicate an abdominal body fat distribution and an increased risk of health problems.² The effect of waist circumference on disease risk in the approximately 10% of adults who have an extreme BMI (≥ 35 kg/m² or < 18.5 kg/m²) is not provided, because the current waist circumference classification does not provide information about disease risk in such adults. Table 2 presents data from a large, population-based cohort study in the United States that provides estimates of disease risk when waist circumference is added to BMI.² For example, among men and women with a normal BMI, an increased waist circumference appears to confer about a 2-fold increase in risk for some diseases. Among those with a BMI of 30–34.9 kg/m², an increased waist circumference con-

Table 1: Body weight classification and risk of health problems by BMI and waist circumference

Measure	Weight classification	Risk of health problems
BMI, kg/m²		
< 18.5	Underweight	Increased
18.5–24.9	Normal weight	Least
25.0–29.9	Overweight	Increased
≥ 30.0	Obese	
30.0–34.9	Class I	High
35.0–39.9	Class II	Very high
≥ 40.0	Class III	Extremely high
Waist circumference		
Men		
< 102 cm (40 in)		Lower
≥ 102 cm (40 in)		Increased
Women		
< 88 cm (35 in)		Lower
≥ 88 cm (35 in)		Increased

PRACTICE


fers a greater than 14-fold increased risk for certain diseases among women but a lesser increase in risk among men. These examples are not necessarily what may be seen in other populations.

Incorporating BMI and waist circumference into an overall risk assessment: The BMI and waist circumference classifications are derived from health risk assessments in large, heterogeneous populations, and consequently

the application of BMI and waist circumference to assess the overall risk of health problems in individual patients will vary from person to person and will depend on the presence of other factors that may influence risk. The BMI and waist circumference should be combined with other determinants of risk (e.g., smoking, concomitant disease, diet, physical activity, and personal and family weight history) to estimate overall disease risk.

For example, physical inactivity is a contributor to all-cause mortality and cardiovascular mortality, irrespective of BMI.³ What may be underappreciated is the importance of BMI and waist circumference compared with the other, more traditional, risk factors. Until recently, obesity was considered to increase the risk of coronary artery disease through its association with hypertension, dyslipidemia and diabetes. However, a BMI of

- Weight and height are required to measure a person's BMI, which is calculated by dividing weight (kilograms) by height (metres squared).
- Weight is measured with the patient wearing light clothing and no footwear and is recorded to the nearest 0.2 kg (1/2 lb).
- Height is measured with the person wearing no shoes, standing in an erect position, looking straight ahead with feet together and heels against a wall or measuring board. A horizontal bar or another similar device should be lowered so that it rests flat on the top of the person's head. Height is recorded to the nearest 0.5 cm (1/4 inch).
- For a quick determination of BMI, use a straight-edge to help locate the point on the chart below where height (in centimetres or inches) and weight (in kilograms or pounds) intersect. Read the number on the dashed line closest to this point. For example, a person who weighs 69 kg and is 173 cm tall has a BMI of about 23.



Note: The BMI cannot be used to distinguish between increased body weight due to adiposity or fluid retention (e.g., peripheral edema), although this distinction should be apparent clinically. The BMI may have limitations in people who are very muscular or very lean, in those who are very tall or very short, in people with long limb lengths relative to trunk size (e.g., those with Marfan syndrome) and in people with short limb lengths relative to trunk size (e.g., those with achondroplasia).

Fig. 1: Measuring body mass index (BMI).

30 kg/m² or greater is an independent risk factor for coronary artery disease, conferring a 2- to 4-fold increase in disease risk, which is comparable to the effect of hypertension on risk of coronary artery disease.⁴

Monitoring BMI and waist circumference over time: The BMI and waist circumference reflect a person's risk status at a single time point. The pattern of BMI and waist circumference over the long term may provide additional information. For example, upward trends in the BMI and waist circumference in people with impaired glucose tolerance may indicate an increased risk of type 2 diabetes. Conversely, downward trends in BMI and waist circumference with unintentional weight loss may indicate an increased risk of disease such as occult cancer.

Weight loss interventions: The need for weight loss interventions should be determined not only on the basis of a person's BMI and waist circumference, but also on the presence or absence of concomitant disease and other patient characteristics. In obese adults with obesity-related diseases such as type 2 diabetes, the goal may be modest weight loss (5%–10% body weight) and lifestyle interventions to improve glycemic and blood pressure control and to reduce the risk of myocardial infarction and stroke. On the other hand, in otherwise healthy obese adults, the weight management goal may be to prevent further weight gain and obesity-related diseases through lifestyle interventions alone.

In what groups may the BMI and waist circumference classification not be applicable?

Elderly people: Interpreting the BMI and waist circumference in elderly patients requires caution. People aged 65 years or older who are overweight may not be at increased risk of

health problems, whereas those who have a BMI in the low end of the normal range (between 18.5 kg/m² and the low 20s) may be at increased risk. Thus, the normal-weight or low-risk BMI range may be higher and wider (e.g., 22–29 kg/m²) for

elderly people than the range for younger adults (18.5–24.9 kg/m²). Weight loss in elderly people, especially loss of fat-free mass, regardless of the baseline BMI, is associated with an increased risk of death. Consequently, screening for weight




Fig. 2: Measuring waist circumference.

Waist circumference	BMI category		
	Normal 18.5–24.9 kg/m ²	Overweight 25–29.9 kg/m ²	Obese class I 30–34.9 kg/m ²
Men: < 102 cm Women: < 88 cm	Least risk	Increased risk	High risk
Men: ≥ 102 cm Women: ≥ 88 cm	Increased risk	High risk	Very high risk

Fig. 3: Applying BMI and waist circumference measurements to assess a patient's risk of health problems.

Table 2: Effect of increased waist circumference* on disease risk when considered in addition to BMI category

Disease	BMI category; adjusted odds ratio (95% confidence interval)†					
	Normal BMI 18.5–24.9 kg/m ²		Overweight BMI 25–29.9 kg/m ²		Obese class I BMI 30–34.9 kg/m ²	
	Women	Men	Women	Men	Women	Men
Hypertension	1.9 (1.3–2.8)	2.3 (0.6–8.7)	2.0 (1.4–3.0)	1.5 (1.1–2.0)	15.9 (4.2–60.3)	1.1 (0.4–3.0)
Type 2 diabetes	1.5 (0.7–3.4)	2.2 (1.1–4.2)	4.1 (1.9–8.7)	2.0 (0.8–5.1)	14.8 (1.7–130.1)	1.0 (0.3–3.2)
Metabolic syndrome	1.8 (1.1–2.9)	0.7 (0.1–3.7)	2.2 (1.2–3.9)	2.1 (1.5–3.2)	28.6 (3.7–220.2)	2.9 (1.3–6.8)

*Increased waist circumference = values of ≥ 102 cm in men and ≥ 88 cm in women. Subjects with waist circumferences below these values were used for the reference group.

†Adjusted for age, race, physical activity, smoking, alcohol intake and poverty-income ratio.

Adapted, with permission, from *Arch Intern Med* 2002;162:2074-9.⁷ Copyright © 2002 American Medical Association.

loss may be a more useful means of determining health risk in elderly people.

Non-white people: Although the body weight classification can be applied to all ethnic groups in Canada, health care providers should be aware of limitations in applying this classification to non-white people. A recent study involving Asian people suggested that BMI cutoffs for overweight and obesity start at 23 kg/m² and 27 kg/m² respectively.⁵ For a given BMI range and waist circumference level, black people appear to be at lower risk of health problems than white people, which suggests the need for higher cutoff levels for BMI and waist circumference to identify increased risk in black populations. In Canadian First Nations and Inuit populations, obesity and type 2 diabetes are major health problems, with over 50% of Aboriginal people having one of these conditions.⁶ Whether the BMI and waist circumference cutoffs used for the general population are appropriate for Aboriginal populations requires additional research.

People with healthy lifestyle habits: The BMI and waist circumference should be applied with caution in people who are physically active or have other

healthy lifestyle habits, because these factors mitigate the risk associated with an increased BMI or waist circumference.

A suggested 3-step approach to using the BMI and waist circumference in clinical practice

1. *Periodic BMI and waist circumference measurement:* Measurement of the BMI and waist circumference should be part of a routine health assessment to screen for overweight and obesity. Measurements should be taken at least every 2 years.
2. *Risk assessment:* The assessment of a patient's risk of health problems should be based on his or her BMI and waist circumference as well as other determinants of risk (e.g., presence or absence of obesity-related disease, lifestyle habits) to establish the person's overall risk profile.
3. *Assessment of need for interventions:* The patient's overall risk profile will, in turn, determine whether lifestyle interventions (e.g., balanced diet, physical activity), weight loss interventions (e.g., dietary therapy) and drug therapy (e.g., antihypertensive or lipid-lowering drugs) are warranted to reduce his or her risk of health problems.

James D. Douketis
St. Joseph's Hospital
Department of Medicine
McMaster University
Hamilton, Ont.

Gilles Paradis
McGill University Health Centre
Montréal, Que.

Heather Keller
Department of Family Relations
and Applied Nutrition
University of Guelph
Guelph, Ont.

Chantal Martineau
Health Canada
Ottawa, Ont.

For the Expert Working Group
on the Canadian Guidelines for Body
Weight Classification in Adults

References

1. Katzmarzyk PT. The Canadian obesity epidemic, 1985–1998. *CMAJ* 2002;166(8):1039-40.
2. Janssen I, Katzmarzyk PT, Ross R. Body mass index, waist circumference, and health risk: evidence in support of current National Institutes of Health guidelines. *Arch Intern Med* 2002;162:2074-9.
3. Hu FB, Willett WC, Li T, Stampfer MJ, Colditz GA, Manson JE. Adiposity as compared with physical activity in predicting mortality among women. *N Engl J Med* 2004;351:2694-703.
4. Eckel RH, Krauss RM; for the AHA Nutrition Committee. American Heart Association call to action: obesity as a major risk factor for coronary heart disease. *Circulation* 1998;97:2099-100.
5. WHO Expert Consultation. Appropriate body-mass index for Asian populations and its implications for policy and intervention strategies. *Lancet* 2004;363:157-63.
6. Young TK, Reading J, Elias B, O'Neil JD. Type 2 diabetes mellitus in Canada's First Nations: status of an epidemic in progress. *CMAJ* 2000;163(5):561-6.